



Contents lists available at ScienceDirect

Fisheries Research

journal homepage: [www.elsevier.com/locate/fishres](http://www.elsevier.com/locate/fishres)



## Genetic population structure of the Swordfish (*Xiphias gladius*) in the southwest Indian Ocean: Sex-biased differentiation, congruency between markers and its incidence in a way of stock assessment

D. Muths<sup>a,\*</sup>, P. Grewe<sup>b</sup>, C. Jean<sup>a</sup>, J. Bourjea<sup>a</sup>

<sup>a</sup> IFREMER Délégation Réunion, Rue Jean Bertho, BP60, 97822 Le Port, France

<sup>b</sup> CSIRO Marine and Atmospheric Research, Castray Esplanade, Hobart, Tasmania 7000, Australia

### ARTICLE INFO

#### Article history:

Received 26 September 2008

Received in revised form 6 March 2009

Accepted 9 March 2009

#### Keywords:

Swordfish

Microsatellites

mtDNA

Population genetic

Indian Ocean

### ABSTRACT

Genetic variation was surveyed at 11 microsatellite loci and at 517 bp of the mitochondrial control region to investigate the presence of genetic stock structure in swordfish (*Xiphias gladius*) in four proximal localities of the southwest Indian Ocean. One aim of this study was to serve as a preliminary examination for congruency of structure detected by these two genetic markers, prior to conducting a more comprehensive basin-wide survey of the Indian Ocean and nearby surrounding areas. Analyses of multilocus microsatellite genotypes and mitochondrial control region sequences both revealed a great homogeneity between samples. Genetic diversity detected at the regional scale was not significantly higher than detected at the local scale. Results suggest that the southwest Indian Ocean globally functions as a unique panmictic population. However, some discrete genetic differences appeared that could possibly indicate influence from a second genetic pool in the northern part of the Indian Ocean. This structure appeared to be sex-dependent with genetic differences higher among female than among male samples. This result may indicate a higher level of spawning area fidelity for females with a subsequent sampling bias tending to homogenise male genotypic distributions.

© 2009 Elsevier B.V. All rights reserved.

### 1. Introduction

The swordfish *Xiphias gladius* is one of the most widely distributed species of pelagic fish, commonly found in the tropical and temperate zones of the Atlantic, Indian and Pacific Oceans. This species is heavily exploited by commercial fisheries worldwide, mainly by drifting longline fisheries. Despite a constant increase of fishery effort, captures of swordfish have decreased since 2000 in the Indian Ocean (Indian Ocean Tuna Commission, 2008). On the basis of the 2008 stock indicators, the IOTC concluded that the recent catch level (about 319,000 t from 2002 to 2006) have been around the current estimate of maximum sustainable yield (31,500 t, 80% confidence limits: 24,500 t–34,400 t) and is so in acceptable limits. While the assessments indicate that the swordfish stock for the whole Indian Ocean is probably not currently overfished, catch rate data from the southwest Indian Ocean suggest that overfishing might occur in localised areas (IOTC, 2008). Consequently, management measures such as quotas introduction may be considered for a sustainable exploitation. However, in the

absence of a clear definition of stock structure, determining the appropriate allocation of the resource will be impossible. In the case of quotas introduction, for example, the application of a unique quota on a mixed population is unfavourable for the species and may lead to stock depletion, whereas the application of several quotas on a unique population may penalize the fishing activity (Avisé, 1998). The artificial spatial scale of stock assessment and management must match with the natural spatial scale of target populations (Francis et al., 2007). In this context, improving knowledge on the population structure of swordfish is the first information needed by managers for defining relevant management measures.

Resolving stock structure in a highly migratory species such as the swordfish presents unique and interesting challenges especially considering observed sexually dimorphic life history strategies. As a general rule, adult female swordfish migrate to temperate areas for feeding during summer, then move to warmer waters for spawning. In contrast males appear to remain more abundant in tropical regions and rarely undertake long-distance migrations observed for females (De Martini et al., 2000; Palko et al., 1981). Tag-recapture experiments showed that even if some swordfish are able to undertake long-distance inter-oceanic movements (at the scale of an ocean; Sedberry and Loefer, 2001; Takahashi et al.,

\* Corresponding author. Tel.: +33 262 554 733.

E-mail address: [Delphine.Muths@ifremer.fr](mailto:Delphine.Muths@ifremer.fr) (D. Muths).

2003), 20% of tagged swordfish are recaptured close to the point of release, revealing important inter-individual plasticity in migrating behaviour and suggesting residency or homing behaviour for some individuals within specific regions (Sedberry and Loefer, 2001). Thus, considering dispersal of the swordfish, it may exist great disparities between the maximal dispersal range (mainly linked with feeding behaviour) and the reproductive effective dispersal range (more important in term of population replenishment and structure).

Genetic studies constitute an efficient mean to determine effective dispersal and delineate stock boundaries (Palumbi, 2003), even for highly migratory species. For example, whereas all tuna species possess similar high migration abilities, differentiation has been detected at various scale: within an ocean basin for bluefin tuna (in Mediterranean; Carlsson et al., 2004), both within and among oceans for the yellowfin tuna (Diaz-Jaimes and Uribe-Alcocer, 2006; Ely et al., 2005; Ward et al., 1997) and for the bigeye tuna (Alvarado Bremer et al., 1998; Durand et al., 2005). On the other hand, no differentiation was observed among oceans for skipjack tuna (Ely et al., 2005). In the case of swordfish, structure appears quite important with populations subdivided on oceanic and infra-oceanic scales, in the Atlantic (Alvarado Bremer et al., 1996, 2005) and Pacific oceans (Lu et al., 2006; Reeb et al., 2000) and in the Mediterranean Sea (Kotoulas et al., 1995). In a general way, swordfish structure has been less studied in the Indian Ocean. Some corridors seem to exist between oceans but these appear to be constrained and delineated by equatorial boundaries. For example, swordfish appear quite similar between South-Indian and South-Atlantic oceans, as well as between South-Indian and Pacific oceans (Chow and Takeyama, 2000; Lu et al., 2006), more similar between neighbouring oceans than from the southern to the northern parts of the same ocean. While most of these genetic studies on the swordfish conducted up until now have involved only one genetic marker, conclusions based on nuclear and mitochondrial DNA data seemed concordant such as observed differentiation in Atlantic Ocean versus the Mediterranean Sea (Alvarado Bremer et al., 2005). However, genetic structure defined in the Indian Ocean with two kinds of markers showed some discrepancies. Indeed, Jean et al. (2006)'s study failed to show population differentiation on the basis of microsatellite data in the southwest Indian Ocean in agreement with what could be suspected for a species displaying a high capacity of migration. On the other hand, Lu et al. (2006) showed with mitochondrial sequences also obtained in the southwest Indian Ocean that gene flow between adjacent populations appeared to be quite reduced or even absent. Even if the uniparental inheritance of mtDNA tends to accentuate genetic differences among population compared to nuclear genes, it does not capture the entire genetic history that is fundamental in the case of defining population structure as required for fisheries management. As such differences in conclusions can have drastic impact on stock assessment, it is imperative to determine whether these differences are due to sampling area, sample size, or to genetic marker discordance.

In order to further examine and clarify whether nuclear and mitochondrial markers are useful and complementary for swordfish stock discrimination, we examined variation at eleven DNA microsatellite loci and compared that with analysis of mitochondrial haplotype data (517-bp of the control region or d-loop). A total of 337 samples was examined from four different sites of the southwest Indian Ocean. The major objective of this study was to provide preliminary support in the form of a pilot study for using both DNA microsatellite and mtDNA markers to further assess population genetic structure of Indian Ocean swordfish in a global context (i.e. the Indian Ocean and its connections with the neighbouring oceanic basins) in an effort to assist better management of this important commercial species.




Fig. 1. Geographic location of the four IOTC fishery statistical square (5° × 5°) wherein *X. gladius* were sampled for this study.

## 2. Materials and methods

### 2.1. Biological materials

Swordfish were sampled from four localities in the southwest Indian Ocean, respectively, in fishery statistical square around the islands of Glorieuses (11°S, 46°E), Seychelles (5°S, 56°E), Reunion (21°S, 56°E) and in the south of Madagascar (31°S, 43°E), respectively, called GLO, SEY, RUN and MADA (Fig. 1). Muscle tissue biopsies from a total of 337 swordfish were collected onboard commercial fishing vessels between February 2005 and May 2006. Tissues were stored in ethanol 90% or in 20% Dimethylsulfoxide (DMSO) saturated salt solution (Dutton, 1996) and frozen until DNA was isolated. All the swordfish sampled were measured (LCK = Length from Cleithrum to Keel, i.e. fish length without head and caudal fin) and sexed (Table 1).

### 2.2. Genetic analysis

Total genomic DNA was extracted using DNAeasy Tissue Kit (Qiagen) following the manufacturers instructions. A 517 base pair (bp) fragment representing a small section of the mitochondrial d-loop of control region was amplified by PCR using the primers defined by Alvarado Bremer (1996; L15998: 5'-TACCCCAAACCTCCCAAAGCTA-3'; H235: 5'-TGAATTAGGAACCCAGATGCCA-3'). Reactions were performed in 25 µl containing 1 × PCR buffer, 2 mM MgCl<sub>2</sub>, 20 µM of each dNTPs, 0.5 µM of each primer, 0.5 U of Advantage Polymerase Taq (Ozyme), 25 ng of genomic DNA. Cycling parameters were 93 °C for 3 min, followed by 35 cycles of 93 °C for 40 s, 60 °C for 50 s, and 72 °C for 40 s and a final elongation at 72 °C for 2 min. PCR products were purified and sequenced on an ABI 3100 sequencer (Macrogen Inc.). Sequences were run forward and reverse. They were checked and edited using Chromas version 1.6 (McCarthy, 1997) and aligned using ClustalW (Thompson et al., 1994) in BioEdit Sequence Alignment Editor (Hall, 1999). Sequences were submitted to GenBank (Accession number EU202452–EU202642).

Eleven microsatellite loci were used, eight from Reeb et al. (2003: Xg-55, Xg-56, XG-66, Xg-75, Xg-144, Xg-166, Xg-379 and Xg-396) and three newly developed loci (D2A, D2B and C8, with D2A and D2B linked loci on the same sequences). For isolation and characterization of these 3 new DNA microsatellite primers, first was total genomic DNA isolated from a single individual and then

**Table 1**  
Characteristics of 3 new microsatellite loci of *Xiphias gladius*. Repeat motif is derived from the sequenced clone.

Locus name	GenBank accession no.	Repeat motif	Primer (5'–3')
D2A	Not yet available. . .	(CTT)6	F-5'-CAGTCGGGGCGTCATCACTCAAAGTAACTTCCAAAGTAACTCT-3'; R-5'-GTTTCACITCCAGCCAAACTCTGTTCG-3'
D2B		(CAGT)8	F-5'-CAGTCGGGGCGTCATCAAAGCAACAACATTGTCTTCTG-3'; R-5'-GTTTCTGGCGTGAACGTGGCTCAATCC-3'
C7		(CTAT)22	F-5'-CAGTCGGGGCGTCATCACTTCAATGTAGAGATGGCAGG-3'; R-5'-GTTTCAAATGTCGGTGGAGCTGTGGACAGA-3'

sent to Genetic Identification Services in California for development of four libraries enriched for sequences containing microsatellite repeats. Each library was enriched for the presence of a specific tetra repeat motif: CAGA, CATC, TAGA or TGAC. Sequences from these four enriched libraries were analysed for the presence of repeats and primers were designed for a few promising loci using Oligo software 6.8 (Molecular Biology Insights, Inc.). Once primer pairs were chosen (Table 1), a CAG tag (5'-CAGTCGGGGCGTCATCA-3'; see Schable et al., 2002) was added to the 5' end of one of each primer pair depending on which gave the least amount of secondary structure. This tag allows the use of a third primer in the PCR (CAG) that is fluorescently labelled for detection on the ABI capillary and gel based genotyping systems (Boutin-Ganache et al., 2001). An additional sequence (GTTTC) was added on the other primer of each pair to promote a-tailing of the fluorescent strand. These 3 microsatellite loci were amplified using AmpliTaq Gold (Applied Biosystems, Inc.) in 50 µl reaction using standard 1.5 mM MgCl<sub>2</sub>. Cycling was kept standard with one hold at 95 °C for 12 min followed by 40 cycles of (94 °C/15 s denature; then 50 °C/30 s anneal; and 72 °C/1 min extension) followed by a final extension at 72 °C for 20 min. The 8 loci from Reeb et al. were amplified using the same PCR reaction for mtDNA, but with Red Gold Star DNA Polymerase (EuroGenTec), and using cycling parameters described in FitzSimmons et al. (1997). Amplified fragments were separated on an ABI Prism 3100 genetic analyser. Alleles were scored using a co-migrating size standard (Genescan500, Applied Biosystems, Inc.) and identified using GeneMapper4 (Applied Biosystems Inc.).

### 2.3. Statistical analyses

#### 2.3.1. Mitochondrial DNA

Genetic variation among mitochondrial sequences was estimated as follows: for each population, the haplotype ( $H_d$ ) and nucleotide ( $\pi$ ) diversities, Tajima's ( $D$ ) statistic and Fu's ( $F$ )  $F$  statistic were examined using the DNAsp 4.0 software (Rozas et al., 2003). Tajima's  $D$  and Fu's  $F$  statistics test for departures from equilibrium

between mutation and drift; significantly negative values indicate population expansion or selective influence. Fu's  $F$  statistics is more sensitive to recent demographic expansion (Fu, 1997). Pair-wise genetic distances ( $\phi_{st}$ ) were estimated between samples using Arlequin 2.000 (Schneider et al., 2001). In all cases, critical significance levels for multiple testing were corrected using a sequential Bonferroni procedure (Rice, 1989). An AMOVA (Analysis of molecular variance; Excoffier et al., 1992) was performed using Arlequin 2.000 (Schneider et al., 2001). DNAsp 4.0 (Rozas et al., 2003) was also used to estimate the nearest neighbour statistic,  $Snn$  (Hudson, 2000).  $Snn$  is a measure of how often the 'nearest neighbours' (in sequence space) are from the same locality (in geographical space).  $Snn$  varies from 0 to 1: under 0.5, it is assumed that populations are in panmixia, and values closer to 1 indicate that populations are differentiated.  $Snn$  is particularly suitable when haplotype diversity is large (Hudson, 2000). Neighbour-joining trees, based on Kimura-2 parameter distance (Kimura, 1980), were constructed using the Mega 2.1 software (Kumar et al., 2001).

#### 2.3.2. Microsatellites

Allele frequencies, genetic diversity for each population, and genetic differentiation between populations were estimated from microsatellites following classical population estimators implemented in the Genepop 3.4 software (Raymond and Rousset, 1995). Values estimated include the mean number of alleles per population ( $N_{all}$ ), and the observed ( $H_{obs}$ ), and expected ( $H_{nb}$ ) heterozygosities (Nei, 1987). In addition, allelic richness ( $R_s$ ) was estimated using Fstat 2.9.3.2 (Goudet, 1995). The null hypothesis of independence between loci was tested from statistical genotypic disequilibrium analyses using Genepop 3.4 (Raymond and Rousset, 1995). Deviations from Hardy–Weinberg equilibrium were examined for each population, at each locus, by calculating Wright's fixation index  $F_{is}$  as estimated by Weir and Cockerham's (1984) using the same software. Departure from Hardy–Weinberg equilibrium was then tested using exact tests. Overall levels of genetic differentiation were analysed by calculating the estimator  $\theta$  of the

**Table 2**  
Main characteristics of the four samples of *X. gladius*.

		GLO	MADA	RUN	SEY
Sampling	$N$	105	100	65	67
	Mean length (cm)	73.7 ± 19.0	90.1 ± 9.4	87.2 ± 21.5	75.2 ± 18.4
	Prop. of females	0.47	0.67	0.13	0.58
Mitochondrial diversity	$h$	90	90	59	60
	$H_d$	0.997	0.997	0.997	0.996
	$\pi$	0.020	0.019	0.021	0.019
	Tajima's $D$	-1.54	-1.45	-1.74	-1.41
	Fu's $F$	-1.52	-1.84	-2.54*	-2.04
Microsatellites diversity	Mean $N_{all}$	17.5	16.0	14.3	15.2
	Mean $R_s$	14.6	13.7	13.3	14.2
	Private $N_{all}$	16	8	3	9
	$H_{nb}$	0.784	0.780	0.776	0.771
	$H_{obs}$	0.720***	0.667***	0.640***	0.658***
	$N_e$ estimate	879	585	277	506

Sampling information are: sample size ( $N$ ), mean length of fish (Length from Cleithrum to Keel in cm, ±Standard Deviation) and proportion of females estimated within each samples. Genetic diversities are given for each markers successively. For mtDNA data, information are: number of haplotypes per population ( $h$ ), haplotype diversity ( $H_d$ ), nucleotide diversity ( $\pi$ ), Tajima's  $D$  and Fu's  $F$  statistics. For microsatellites, information are: mean number of alleles ( $N_{all}$ ), allelic richness ( $R_s$  as estimated for a common minimal sample size of 44 individuals), number of private alleles, unbiased ( $H_{nb}$ ) and observed ( $H_{obs}$ ) heterozygosities (Nei, 1987) and effective size estimates ( $N_e$ ). Significant values are noticed by \* $p < 0.05$ ; \*\*\* $p < 0.001$ .

Wright's  $F_{ST}$  Statistic (Weir and Cockerham, 1984) for each locus, and differentiation was then tested using exact tests for the null hypothesis of identity of allelic distributions across populations. Effective population size ( $N_e$ ) were estimated using changes in microsatellite allele frequencies with the software programme NeEstimator (Peel et al., 2004). This software gave point estimation of  $N_e$  using linkage/gametic disequilibrium (Hill, 1981). NeEstimator was not used to estimate the actual long-term inbreeding effective population size but to compare  $N_e$  estimates as relative effective population sizes between samples. AMOVA analysis were performed using Arlequin 2.000 (Schneider et al., 2001). A correspondence factorial analysis was performed on genotype frequencies with the Genetix 4.0 software (Belkhir et al., 2000). To determine if the samples belonged to one or more populations, data were also analysed using the software Structure (Pritchard et al., 2000) which uses iterative computation process to infer the most likely number of populations ( $K$ ) represented in the total sample. For this analysis, an admixture model assuming independent allele frequencies was used and three replicates were run (each with  $1 \times 10^5$  burn-ins and  $5 \times 10^5$  iterations) at  $K$  values from 1 to 4.

### 3. Results

#### 3.1. Genetic diversity and demographic stability

##### 3.1.1. Mitochondrial DNA

A total of 117 variable sites, constituting 240 haplotypes was detected among the mtDNA control region sequences (517 bp) for the 337 swordfish sequenced. Compiling with Reeb et al.' (2000)s dataset previously published in GenBank (Accession number AF199616–AF200183), it appears that all these haplotypes belong to clade I with a divergence intra-clade lower than 1% (compared to a divergence between clades higher than 2.5%).

A similar high level of genetic diversity was encountered in each locality (Table 2). Mean haplotype diversity ( $H_d$ ) and mean nucleotide diversity ( $\pi$ ) were of the same order of magnitude between localities, with  $H_d$  near from 1, and  $\pi$  near from 0.02. Tajima's  $D$  and Fu's  $F$  values were negative but not significant ( $D = -1.6$  and  $F = -2$ ), except for Fu's statistics in RUN locality.

##### 3.1.2. Microsatellites

Mean number of alleles and allelic richness were of the same order between the four localities respectively varying from 14.3 to 17.5 and from 13.3 to 14.6 (Table 1), with each time the lowest value in RUN and the highest value in GLO. The GLO population had the highest number of private alleles at 16 versus a range of 3–9 in the three other localities).  $N_e$  estimates varied from a low of 277 in RUN to 879 in GLO (Table 2). No loci were in disequilibrium ( $p < 0.001$ ) over the whole dataset, supporting the independent assortment of alleles at different loci. Heterozygote deficiencies were highly significant in all samples with values ranging from a low of 0.079 for GLO to 0.176 for RUN (Tables 2 and 3).

#### 3.2. Population structure

##### 3.2.1. Mitochondrial DNA

Pairwise  $\phi_{ST}$  estimates between localities are presented in Table 4. Mean value of  $\phi_{ST}$  was weak ( $=0.01$ ) with only one of the six values significant, between SEY and RUN samples. The two highest  $\phi_{ST}$  values were observed in the SEY sample. Interestingly, an AMOVA across the four samples demonstrated a small and non-significant  $\Phi_{ST}$  value (0.005,  $p = 0.15$ ). When hierarchical AMOVA analysis were undertaken with grouping schemes in agreement with significant pairwise  $\phi_{ST}$  estimates on the localities (*i.e.* SEY sample isolated from the three others or RUN sample isolated from the three others), more than 99% of the variance was

**Table 3**

Mono- and multi-loci estimates of the fixation index  $F_{IS}$  within each locality of *X. gladius*. Tests of significance were performed with Genetix 4.0 (Belkhir et al., 2000). Allele size range (in base pairs) and number of alleles per locus are also given.

Locus	Size range	Nall	Monolocus $F_{IS}$			
			GLO	MADA	RUN	SEY
X55	79–191	47	0.149***	0.330***	0.405***	0.170***
X56	115–159	20	0.025	0.110***	0.064	0.094*
X66	110–140	12	0.094*	0.115***	0.150***	0.080
X75	142–276	53	0.031	0.332***	0.343***	0.182***
X144	151–172	8	–0.005	0.033	0.076	0.140
X166	120–144	9	–0.007	0.079	0.078	0.048
X379	100–142	15	0.020	–0.013	0.099	0.127*
X396	107–137	9	0.182***	0.326***	0.381***	0.300***
D2A	287–296	4	0.079	0.048	0.064	0.230***
D2B	142–202	16	0.044	–0.007	0.019	0.060
C8	136–240	28	0.223	0.131	0.164	0.221
Multiloci average $F_{IS}$						
			0.079***	0.146***	0.176***	0.148***

\*  $p < 0.05$ .

\*\*\*  $p < 0.001$ .

observed within the samples ( $\Phi_{ST} < 0.002$ ,  $p > 0.05$ ) with a non-significant variance associated with the partition in two groups ( $\Phi_{SC} < 0.001$ ,  $p > 0.05$ ). To further test samples homogeneity, the nearest neighbour statistic ( $Snn$ ) was calculated on the mtDNA control region sequences. The test revealed a non-significant association between sequence similarity and geographical location ( $Snn = 0.288$ ,  $p = 0.13$ ). A neighbour-joining tree based on average pairwise distances estimated from the 517-bp mtDNA sequences between samples is presented in Fig. 2. Samples from the four localities appeared well mixed. Adding our sequences to Lu et al. (2006)'s Indian Ocean sequences previously published in GenBank (Accession number DQ076502–DQ076643), pairwise  $\phi_{ST}$  estimates revealed no more differentiation. This could be due to the low sampling size of Lu et al. (2006)'s samples (four times lower than ours) or to shorter length of our sequences (517 bp against 819 bp).

##### 3.3. Microsatellites

Pairwise multilocus  $\theta$  values between localities are presented in Table 4. Mean value of  $\theta$  was weak ( $=0.02$ ) with all the non-null  $\theta$  values involving either SEY or GLO samples. Only one significant  $\theta$  value was observed among six sample site comparisons. This was between SEY and GLO and was likely due to significant differentiation observed at 5 of the 11 loci tested. An AMOVA across the four samples demonstrated a small and non-significant  $\Phi_{ST}$  value (0.000,  $p = 0.510$ ). When hierarchical AMOVA analysis were undertaken with grouping schemes in agreement with significant pairwise  $\theta$  values between localities (*i.e.* SEY or GLO sample isolated from the three others), more than 99% of the variance was observed within the samples ( $\Phi_{ST} < 0.001$ ,  $p > 0.05$ ) with a non-


**Table 4**

Pairwise values of genetic differentiation in *X. gladius* localities. Pairwise  $\phi_{ST}$  values obtained from the mtDNA sequences dataset are above the diagonal with tests of significance performed with Arlequin 2.0 (Schneider et al., 2001). Multiloci  $\theta$  values obtained from the microsatellite dataset are below the diagonal with tests of significance performed with Genetix 4.0 (Belkhir et al., 2000).

	GLO	MADA	RUN	SEY
GLO				
MADA	0.001	0.000	0.000	0.001
RUN	0.000	0.000	0.001	0.001
SEY	0.005***	0.002	0.003	0.003*


\*  $p < 0.05$ .

\*\*\*  $p < 0.001$ .



**Fig. 2.** Neighbour-joining tree based on pairwise number of differences between haplotypes of *X. gladius* from the four localities of the southwest of Indian Ocean. Samples are respectively represented by black squares for GLO, grey circles for Mada, white squares for RUN and white triangles for SEY.

significant variance associated with the partition in two groups ( $\Phi_{SC} < 0.001, p > 0.05$ ). The analysis made with Structure suggested that the highest likelihood of obtaining such data was to consider that only one population ( $K = 1$ ) existed. The likelihood decreased when estimates were made with one population to two (over three independent simulations:  $\ln P(D)$  for  $K = 1$  and  $K = 2$  were  $-14217$  and  $-14536$ , respectively) providing some evidence against subdivision. A correspondence factorial analysis was performed on genotype frequencies. Results of this multivariate analysis are presented in Fig. 3; 81% of genetic variance was synthesized by the two first axis (respectively 39.3% and 31.7% by axis 1 and axis 2). Some groupings seem to exist in accordance with sampling locality, mainly due to the segregation of SEY against GLO on the first axis, whereas these two samples appeared opposed to MADA and RUN on the second axis. However the distribution of individuals along the axes showed a great disparity and a high degree of superposition.



**Fig. 3.** Correspondence factorial analysis, performed on genotype frequencies of *X. gladius*. Each point represents a given individual whose symbol corresponds to its sampling locality.

**Table 5**

Pairwise microsatellite multiloci  $\theta$  values of genetic differentiation between males and females swordfish with tests of significance performed with Genetix 4.0 (Belkhir et al., 2000). Differentiation between females of two localities are below the diagonal, between males above the diagonal and between males and females from a same locality on the diagonal. Samples size are noticed into parentheses.

Males	GLO (57)	MADA (15)	RUN (31)	SEY (26)
Females				
GLO(49)	0.005***	0.005	0.005*	0.001
MADA(33)	0.005	0.000	0.006	0.000
RUN (6)	0.009	0.000	0.010	0.005
SEY(25)	0.013***	0.004	0.006	0.003

\*  $p < 0.05$ .  
\*\*\*  $p < 0.001$ .

**3.4. Sex-dependent analysis**

As sex identification was available for each swordfish, some statistics were re-estimated, for males and females, of each locality or of all localities. Levels of genetic diversities of both markers were of the same order within each sex category. Allelic richness was however always higher for female than for male but not in a significant way. Pairwise  $\phi_{st}$  estimates between sex categories within each locality were weak. The previous significant value found between SEY and RUN disappeared (probably because of the very small number of females in RUN locality ( $n = 6$ )). Notably, the previous pairwise multilocus  $\theta$  value found between SEY and GLO was still significant indicating that significant differences were only due to females ( $\theta = 0.01, p < 0.001$ ), as males from SEY and GLO showed no significant differences (see Table 5). Other pairwise differentiation involving females from GLO were significant (i.e. with males from GLO or with females from MADA; see Table 5). In a global way, all pairwise differentiation values involving females were higher than the same values estimated between males. Two hierarchical AMOVA (one on mtDNA data, the second on microsatellites) were undertaken with partitioning in two sex groups. More than 99% of the variance was observed within the samples ( $\Phi_{ST} < 0.001, p > 0.05$ ) with a non-significant variance associated with the partition of the two sex groups ( $\Phi_{SC} < 0.001, p > 0.05$ ).

**4. Discussion**

This study aimed to yield results on two different levels. The first aim was to evaluate genetic differences between swordfish sampled in four distinct localities of the southwest Indian Ocean (SWIO). An other aim was to examine the amount of congruency between the two genetic marker types used to examine the population structure and its implication in the case of a regional swordfish multi-stock assessment program.

**4.1. Genetic structure of the swordfish in SWIO**

The analysis of mitochondrial sequences of *X. gladius* has revealed a very high level of mitochondrial diversity (nearly all individual displayed a distinct haplotype) as well as a high microsatellite polymorphism (some loci showing up to 40 alleles). Both markers showed a great genetic homogeneity between the four samples. *Snn* statistic as well as results obtained with the software Structure pointed out the existence of a unique pool of genes. In the same way, analysis of molecular variance mainly failed to find a genetic structure among the four localities sampled in the SWIO. The very high levels of genetic diversity and the lack of differentiation detected at a large spatial scale (i.e. about 2000 miles from North to South of the sampling area) is thus well in agreement with the assumption of a large population size that is not very sensitive to genetic drift (De Woody and Avise, 2000). This last

point agrees with Tajima's  $D$  and Fu's  $F$  statistics that indicate these samples appear demographically stable. The hypothesis of a large effective population size that do not fluctuate greatly over time is easily understandable for swordfish with fecundities of several millions of eggs per female (Palko et al., 1981). Such a finding is also in agreement with the hypothesis of long range dispersal typical of large pelagic fishes (Waples, 1998).

Results obtained from both mtDNA and microsatellites are consistent with the idea that swordfish of the SWIO belong to a single unique panmictic population. However, some weak differentiation seems to exist within the SWIO, and thus despite the important within-sample diversity that has been noticed to considerably reduce the ability of detecting between-samples structure, either on mtDNA data (Charlesworth, 1998) or on microsatellites studies (Hedrick, 1999; O'Reilly et al., 2004). Jean et al. (2006)'s study failed to demonstrate genetic structure of swordfish within the SWIO; increasing the sample size and the number of microsatellites loci (three times more samples, two times more microsatellites, between Jean et al. (2006)'s study and our) has thus permit to display heterogeneity within swordfish of the SWIO. The punctual estimation of effective population size (that is the spawning proportion of adults) tends to show higher values in the two northern localities (GLO and SEY), in parallel with higher allelic richness. This could be in agreement with the hypothesis of a reproductive aggregation in this northern zone (Mejuto et al., 2006) whereas the two southern areas might better represent transition zones between feeding and spawning areas or feeding regions. Some swordfish collected in these northern localities (in GLO or SEY) could thus belong to a second genetic pool. The significant heterozygote deficits observed in all localities could partly be the signature of a Wahlund effect and could confirm the, more or less pronounced, influence of a second genetic pool. This is also in agreement with Lu et al. (2006)'s study which showed the existence of differences between swordfish sampled in the Indian Ocean (in this case, between north of Madagascar and other northern sampling sites). Swordfish within a locality may thus be a mixture from one dominant genetic pool and a second less influent one, with various level of homogeneity depending on the swordfish behaviour, and thus explaining the weak structure observed. These results contrast with those obtained in Atlantic (Alvarado Bremer et al., 2005) that support the homogeneity between spawning and feeding grounds, either in North and South Atlantic.

Moreover, examination of these populations indicates that there are some sex-biased intra-sample differences, that could also explain heterozygote deficits. For example, males in GLO differ from females of the same area and the genetic differences found between SEY and GLO was in fact only due to females. This species is known to migrate to temperate areas for feeding and then move to warmer waters to reproduce (Palko et al., 1981): differences in the level of genetic structure observed when only males or females are involved could thus be the signature of a more pronounced homing behaviour depending on sex (Keeney et al., 2005; Lee et al., 2007). In the present case, higher values for females indicate that they might present a higher fidelity to their reproduction areas than males (Prugnolle and de Meeus, 2002). A lower fidelity by males, means that the male fish are supposed to swim around more and are caught randomly with less structure when they are caught. This is supposed to homogenise the nuclear genetic signal for both sexes, and so the differentiation between females might be higher with the mtDNA. Actually it is not the case. So it could mean that a sampling biased exist, maybe without enough homogeneity in sampling periods. The sampling is not yet sufficient to deal further with sex-biased differences but open a very interesting perspective of research. Conclusions based solely on genetic data would be an insufficient method of defining population structure, without the light of some basic biological informa-

tional framework based on reproductive, feeding, and migrating strategies.

#### 4.2. Congruency of molecular markers in a perspective of stock assessment

Many genetic studies have shown discrepancies in conclusions when using different markers types (Lemaire et al., 2000; Nielsen et al., 2006; Pogson et al., 1995). In such cases, result disparities may partly be explained by differences in mutation rates and in sensitivity to forces that promote population differentiation. Using two genetic markers was our initial option to limit errors, as 'drawing conclusions from single genealogies can be problematic as each is only a single point in the space of all possible genealogies' (Wakeley, 2003). In the present study, both markers seemed to support similar conclusions in the pattern of genetic differentiation. Despite the high genetic diversity levels and the consequent limits of interpretation both markers indeed showed a global pattern of panmixia within a unique population at the scale of the SWIO influenced by a putative second differentiated population in the equatorial region of the Indian Ocean. The putative existence of a second pool in a restricted area of the SWIO is important as this region is supposed to be more affected by stock declines due to high fishing pressures (IOTC, 2008). The existence of a distinct stock have first to be confirmed; but, if it is really the case, genetic data will provide a key information in the aim of swordfish stock assessment in Indian Ocean as decision have to be taken separately for each stock. This encourages further research. Sampling a more extensive area (within the Indian Ocean and neighbouring oceanic basins) may permit estimates of the number of independent gene pools of swordfish, their geographic boundaries, and the level of exchanges between them. The existence of a second putative spawning aggregation in the western part of the Indian Ocean as well as the specificity of bias due to dimorphic population patterns observed in the different sexes also warrants further investigation. Finally, a pertinent strategy for dealing with veracity of conclusions requires a good temporal sampling strategy that is replicated samples over a 2–3 years time frame to fully describe population dynamics for species like the swordfish that has the migratory potential to demonstrate high levels of gene flow (Waples, 1998). Temporal samples will also permit effective estimates of population size with higher precision (Fraser et al., 2007) and help to challenge questions about overfishing.

The present study permits analysis of genetic structure observed in *X. gladius* in the SWIO and to address conclusions of previous studies (Jean et al., 2006; Lu et al., 2006). Our results also underline the necessity of compiling biological data to genetic information to examine swordfish stock structure. Stock estimates clearly require further investigations by increasing the spatio-temporal sampling, both within Indian Ocean and with the neighbouring oceanic basins. As underlined by Francis et al. (2007) in the fourth of their Ten commandments for ecosystem-based fisheries scientists, 'continuing to rely on traditional stock assessments that either ignore or artificially delineate the true spatial structure of fish populations is clearly a recipe for disaster'. A project of this scale would provide much insight and constitute an excellent database for the CTOI and management authorities for both present and future management of swordfish fishery in the Indian Ocean.

#### Acknowledgements

We would like to thank L. Gagnevin and K. Vital (CIRAD-3P) for their help in the laboratory work, P. Bach (Institut de Recherche pour le Développement), Alicia Delgado (Instituto Español de Oceanografía) and the Seychelles Fishing Authorities for their help

in collecting effort. Funding for this project was provided by program DEMOSTEM-STRADA IFREMER.

## References

- Alvarado Bremer, J., Mejuto, J., Gomez-Marquez, J., Boan, F., Carpintero, P., Rodriguez, J., Vinas, J., Greig, T., Ely, B., 2005. Hierarchical analyses of genetic variation of samples from breeding and feeding grounds confirm the genetic partitioning of northwest Atlantic and South Atlantic populations of swordfish (*Xiphias gladius* L.). *J. Exp. Mar. Biol. Ecol.* 327, 167–182.
- Alvarado Bremer, J., Mejuto, J., Grei, T., Ely, B., 1996. Global population structure of the swordfish (*Xiphias gladius*) as revealed by analysis of the mitochondrial DNA control region. *J. Exp. Mar. Biol. Ecol.* 97, 295–310.
- Alvarado Bremer, J.R., Stequert, B., Robertson, N.W., Ely, B., 1998. Genetic evidence for inter-oceanic subdivision of bigeye tuna (*Thunnus obesus*) populations. *Mar. Biol.* 132, 547–557.
- Avise, J.C., 1998. Conservation genetics in the marine realm. *J. Hered.* 89, 377–382.
- Belkhir, K., Borsa, P., Goudet, J., Chikhi, L., Bonhomme, F., 2000. GENETIX 4.02, logiciel sous WindowsTM pour la génétique des populations. Laboratoire Génome, Populations. Interactions CNRS UMR 5000. Université de Montpellier II, Montpellier (France).
- Boutin-Ganache, I., Raposo, M., Raymond, M., Deschepper, C., 2001. M13-tailed primers improve the readability and usability of microsatellite analyses performed with two different allele-sizing methods. *Biotechniques* 31, 24–28.
- Carlsson, J., McDowell, J.R., Diaz-Jaimes, P., Carlsson, J.E.L., Boles, S.B., Gold, J.R., Graves, J.E., 2004. Microsatellite and mitochondrial DNA analyses of Atlantic bluefin tuna (*Thunnus thynnus*) population structure in the Mediterranean Sea. *Mol. Ecol.* 13, 3345–3356.
- Charlesworth, B., 1998. Measures of divergence between populations and the effect of forces that reduce variability. *Mol. Biol. Evol.* 15, 538–543.
- Chow, S., Takeyama, H., 2000. Nuclear and mitochondrial DNA analyses reveal four genetically separated breeding units of the swordfish. *J. Fish Biol.* 56, 1087–1098.
- De Martini, E., Uchiyama, J., Williams, H., 2000. Sexual maturity, sex ratio, and size composition of swordfish, *Xiphias gladius*, caught by the Hawaii-based pelagic longline fishery. *Fish. Bull.* 98, 489–506.
- De Woody, J., Avise, J., 2000. Microsatellite variation in marine, freshwater, and andromous fishes compared with other animals. *J. Fish Biol.* 56, 461–473.
- Diaz-Jaimes, P., Uribe-Alcocer, M., 2006. Spatial differentiation in the eastern Pacific yellowfin tuna revealed by microsatellite variation. *Fish. Sci.* 72, 590–596.
- Durand, J., Collet, A., Chow, S., Guinand, B., Borsa, P., 2005. Nuclear and mitochondrial DNA markers indicate unidirectional gene flow of Indo-Pacific to Atlantic bigeye tuna (*Thunnus obesus*) populations, and their admixture off southern Africa. *Mar. Biol.* 147, 313–322.
- Dutton, P., 1996. Methods for collection and preservation of samples for sea turtle genetic studies. In: Memorandum, N.T.S. (Ed.), Proceedings of the International Symposium on Sea Turtle Conservation Genetics, pp. 17–24.
- Ely, B., Vinas, J., Alvarado Bremer, J.R., Black, D., Lucas, L., Covello, K., Labrie, A.V., Thelen, E., 2005. Consequences of the historical demography on the global population structure of two highly migratory cosmopolitan marine fishes: the yellowfin tuna (*Thunnus albacares*) and the skipjack tuna (*Katsuwonus pelamis*). *Evol. Biol.* 5, 19.
- Excoffier, L., Smouse, P., Quattro, J., 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 131, 479–491.
- FitzSimmons, N.N., Limpus, C.J., Norman, J.A., Goldizen, A.R., Miller, J.D., Moritz, C., 1997. Philopatry of male marine turtles inferred from mitochondrial DNA markers. *Proc. Natl. Acad. Sci. U.S.A.* 94, 8912–8917.
- Francis, R., Hixon, M., Clarke, M., Murawski, S., Ralston, S., 2007. Ten commandments for ecosystem-based fisheries scientists. *Fisheries* 32, 217–233.
- Fraser, D.J., Hansen, M.M., Ostergaard, S., Tessier, N., Legault, M., Bernatchez, L., 2007. Comparative estimation of effective population sizes and temporal gene flow in two contrasting population systems. *Mol. Ecol.* 16, 3866–3889.
- Fu, Y., 1997. Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics* 147, 915–925.
- Goudet, J., 1995. FSTAT (vers. 1. 2): a computer program to calculate *F*-statistics. *J. Hered.* 86, 485–486.
- Hall, T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acid Symp. Series* 41, 95–98.
- Hedrick, P.H., 1999. Perspective: highly variable loci and their interpretation in evolution and conservation. *Evolution* 53, 313–318.
- Hill, W.G., 1981. Estimation of effective population size from data on linkage disequilibrium. *Gen. Res.* 38, 209–216.
- Hudson, R.R., 2000. A new statistic for detecting genetic differentiation. *Genetics* 155, 2011–2014.
- IOTC, 2008. IOTC. Report of the Eleventh Session of the Scientific Committee (1–5 December, 2008). IOTC (Indian Ocean Tuna Commission), Victoria, Seychelles, p. 166.
- Jean, C., Bourjea, J., Jouen, E., Taquet, M., 2006. Stock structure of the swordfish (*Xiphias gladius*) in the Southwest Indian Ocean: a preliminary study. *Bull. Mar. Sci.* 79, 521–526.
- Keeney, D.B., Heupel, M.R., Hueter, R.E., Heist, E.J., 2005. Microsatellite and mitochondrial DNA analyses of the genetic structure of blacktip shark (*Carcharhinus limbatus*) nurseries in the northwestern Atlantic, Gulf of Mexico, and Caribbean Sea. *Mol. Ecol.* 14, 1911–1923.
- Kimura, M., 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* 16, 111–120.
- Kotoulas, G., Magoulas, A., Tsimenides, N., Zouros, E., 1995. Marked mitochondrial DNA differences between Mediterranean and Atlantic populations of the swordfish, *Xiphias gladius*. *Mol. Ecol.* 4, 473–481.
- Kumar, S., Tamura, K., Jakobsen, I., Nei, M., 2001. MEGA2: Molecular Evolutionary Genetics Analysis software. *Bioinformatics* 17, 1244–1245.
- Lee, P.L., Luschi, P., Hays, G.C., 2007. Detecting female precise natal philopatry in green turtles using assignment methods. *Mol. Ecol.* 16, 61–74.
- Lemaire, C., Allegrucci, G., Naciri, M., Bahri-Sfar, L., Kara, H., Bonhomme, F., 2000. Do discrepancies between microsatellite and allozyme variation reveal differential selection between sea and lagoon in the sea bass (*Dicentrarchus labrax*)? *Mol. Ecol.* 9, 457.
- Lu, C., Chen, C., Hui, C., Tzen, T., Yeh, S., 2006. Population genetic structure of the swordfish, *Xiphias gladius* (Linnaeus, 1758) in the Indian Ocean and the West Pacific inferred from the complete DNA sequence of the mitochondrial control region. *Zool. Stud.* 45, 269–279.
- McCarthy, C., 1997. Chromas, Version 1. 41. Griffith University, Brisbane.
- Mejuto, J., Garcia-Cortes, B., Ramos-Cortelle, A., 2006. An overview of research activities on swordfish (*Xiphias gladius*) and the by-catch species caught by the Spanish longline fleet in the Indian Ocean. IOTC.
- Nei, M., 1987. *Molecular Evolutionary Genetics*. Columbia University Press, New York.
- Nielsen, E.E., Hansen, M.M., Meldrup, D., 2006. Evidence of microsatellite hitch-hiking selection in Atlantic cod (*Gadus morhua* L.): implications for inferring population structure in nonmodel organisms. *Mol. Ecol.* 15, 3219–3229.
- O'Reilly, P.T., Canino, M.F., Bailey, K.M., Bentzen, P., 2004. Inverse relationship between *F* and microsatellite polymorphism in the marine fish, walleye pollock (*Theragra chalcogramma*): implications for resolving weak population structure. *Mol. Ecol.* 13, 1799–1814.
- Palko, B., Beardslay, G., Richards, W., 1981. Synopsis of the biology of the swordfish, *Xiphias gladius* Linnaeus. NOAA Technical Report NMFS Circular 441 FAO Fisheries synopsis, 2–15.
- Palumbi, S.R., 2003. Population genetics, demographic connectivity and the design of marine protected areas. *Ecol. Appl.* 13, 146–158.
- Peel, D., Ovenden, J.R., Peel, S.L., 2004. NeEstimator: Software for Estimating Effective Population Size. Molecular Fisheries Laboratory, Queensland Department of Primary Industries, Australia.
- Pogson, G.H., Mesa, K.A., Boullier, R.G., 1995. Genetic population structure and gene flow in the Atlantic Cod *Gadus morhua*: a comparison of allozyme and nuclear RFLP loci. *Genetics* 139, 375–385.
- Pritchard, J.K., Stephens, M., Donnelly, P., 2000. Inference of population structure using multilocus genotype data. *Genetics* 155, 945–959.
- Prugnolle, F., de Meeus, T., 2002. Inferring sex-biased dispersal from population genetic tools: a review. *Heredity* 88, 161–165.
- Raymond, M., Rousset, F., 1995. GENEPOP (version 1. 2): a population genetics software for exact tests and ecumenicism. *J. Hered.* 86, 248–249.
- Reeb, C., Arcangeli, L., Block, B., 2000. Structure and migration corridors in Pacific populations of the swordfish *Xiphias gladius*, as inferred through analysis of mitochondrial DNA. *Mar. Biol.* 136, 1123–1131.
- Reeb, C.A., Arcangeli, L., Block, B.A., 2003. Development of 11 microsatellite loci for population studies in the swordfish, *Xiphias gladius* (Teleostei: Scombridae). *Mol. Ecol. Notes* 3, 147–169.
- Rice, W., 1989. Analyzing tables of statistical tests. *Evolution* 43, 223–225.
- Rozas, J., Sanchez-DelBarrio, J.C., Messeguer, X., Rozas, R., 2003. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* 19, 2496–2497.
- Schable, N., Fischer, R., Glenn, T., 2002. Tetranucleotide microsatellite DNA loci from the dollar sunfish (*Lepomis marginatus*). *Mol. Ecol. Notes* 2, 509–511.
- Schneider, S., Roessli, D., Excoffier, L., 2001. Arlequin (v 2. 0): Software for population Genetics Data Analysis. Genetics and Biometry Laboratory, Department of Anthropology, University of Geneva, Geneva.
- Sedberry, G., Loefer, J., 2001. Satellite telemetry tracking of swordfish, *Xiphias gladius*, off the eastern United States. *Mar. Biol.* 139, 355–360.
- Tajima, F., 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123, 585–595.
- Takahashi, M., Okamura, H., Yokawa, K., Okazaki, M., 2003. Swimming behaviour and migration of a swordfish recorded by an archival tag. *Mar. Freshwater Res.* 54, 527–534.
- Thompson, J.D., Higgins, D.G., Gibson, T.J., 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22, 4673–4680.
- Wakeley, J., 2003. Inferences about the structure and history of populations: coalescents and intraspecific phylogeography. In: Singh, R., Uyenoyama, M. (Eds.), *The Evolution of Population Biology*. Cambridge University Press, Cambridge, pp. 193–215.
- Waples, R.S., 1998. Separating the wheat from the chaff: patterns of genetic differentiation in high gene flow species. *J. Hered.* 89, 438–450.
- Ward, R., Elliott, N., Innes, B., Smolenski, A., Grewe, P., 1997. Global population structure of yellowfin tuna, *Thunnus albacares*, inferred from allozyme and mitochondrial DNA variation. *Fish. Bull.* 95, 566–575.
- Weir, B.S., Cockerham, C.C., 1984. Estimating *F*-statistics for the analysis of population structure. *Evolution* 38, 1358–1370.